

Improper Integrals

Name:

Block:

Seat:

Improper Integrals are integrals where:

- One or both limits of integration are $\pm\infty$
(sometimes called “Type I” or “horizontal” improper integrals)
- The integrand is undefined at one or both of the limits of integration or in between the limits of integration
(the integrand has a vertical asymptote, sometimes called “Type II” or “vertical” improper integrals)

Improper integrals are said to be

- **convergent** if the limit is finite and that limit is the value of the improper integral.
- **divergent** if the limit does not exist.

Examples

$$1. \int_0^{\infty} e^{-x} dx = \lim_{b \rightarrow \infty} \int_0^b e^{-x} dx = \lim_{b \rightarrow \infty} [-e^{-x}]_0^b = \lim_{b \rightarrow \infty} \frac{-1}{e^b} + e^0 = 0 + 1 = 1$$

$$2. \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

$$\text{First we consider } \int_{-\infty}^0 \frac{1}{1+x^2} dx = \lim_{b \rightarrow -\infty} [\arctan x]_b^0 = \arctan 0 - \lim_{b \rightarrow -\infty} \arctan b = 0 - \frac{\pi}{2} = -\frac{\pi}{2}$$

$$\text{Similarly } \int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{b \rightarrow \infty} [\arctan x]_0^b = \lim_{b \rightarrow \infty} \arctan b - \arctan 0 = \frac{\pi}{2} - 0 = \frac{\pi}{2}$$

$$\text{Putting it together, } \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = -\frac{\pi}{2} + \frac{\pi}{2} = \pi$$

p -Test for Integrals Thm 8.7 (page 578):

If $p > 1$, then $\int_1^{\infty} \frac{1}{x^p} dx$ converges to $\frac{1}{p-1}$

If $p \leq 1$, then $\int_1^{\infty} \frac{1}{x^p} dx$ diverges

Examples

1. $\int_1^{\infty} \frac{1}{x^{3/2}} dx$

Following usual methods:

$$\lim_{b \rightarrow \infty} \int_1^b \frac{1}{x^{3/2}} dx = \lim_{b \rightarrow \infty} \int_1^b x^{-3/2} dx = \lim_{b \rightarrow \infty} \left[\frac{1}{-\frac{1}{2}} x^{-1/2} \right]_1^b = \lim_{b \rightarrow \infty} - \left[\frac{2}{x^{1/2}} \right]_1^b = 0 - (-2) = 2$$

But following the theorem:

$$p = \frac{3}{2} > 1, \text{ hence } \frac{1}{p-1} = \frac{1}{\frac{3}{2}-1} = \frac{1}{\frac{1}{2}} = 2$$

2. $\int_1^{\infty} \frac{1}{x^{1/2}} dx$

Following usual methods:

$$\lim_{b \rightarrow \infty} \int_1^b \frac{1}{x^{1/2}} dx = \lim_{b \rightarrow \infty} [2\sqrt{x}]_1^b = \lim_{b \rightarrow \infty} 2\sqrt{b} - 2 = \infty$$

But following the theorem:

$$p = \frac{1}{2} \leq 1, \text{ hence it diverges}$$

Exercises

1. $\int_1^{\infty} \frac{1}{x} dx$ (try it both ways!)

2. $\int_1^{\infty} \frac{1}{x^2} dx$

3. $\int_0^{\infty} \frac{1}{x^2 + 9} dx$

Hint: $\frac{1}{x^2 + 9} = \frac{\frac{1}{9}}{\frac{x^2}{9} + 1}$ reminds me of arctan (see p. 520)

4. $\int_0^1 \frac{1}{\sqrt{x}} dx$

Hint: Use the limit as $b \rightarrow 0^+$

5. $\int_{-1}^1 \frac{1}{x^{2/3}} dx$

6. $\int_{-2}^1 \frac{1}{x^2} dx$

7. $\int_0^1 x \ln x \, dx$

8. $\int_0^\infty \cos x \, dx$

Answers: 1) $\frac{\pi}{8}$, 2) $\frac{\pi}{8}$, 3) $\frac{\pi}{8}$, 4) $\frac{\pi}{8}$, 5) $\frac{\pi}{8}$, 6) $\frac{\pi}{8}$, 7) $\frac{\pi}{8}$, 8) $\frac{\pi}{8}$